

Nitrous oxide emissions pose an increasing climate threat, study finds

Date:

October 7, 2020

Source:


University of East Anglia

Summary:

Rising nitrous oxide emissions are jeopardizing the climate goals of the Paris Agreement, according to a major new study. The growing use of nitrogen fertilizers in the production of food worldwide is increasing atmospheric concentrations of nitrous oxide -- a greenhouse gas 300 times more potent than carbon dioxide that remains in the atmosphere for more than 100 years.

Share:

FULL STORY



Tractor spraying field (stock image).
Credit: © eleonimages / stock.adobe.com

Rising nitrous oxide (N₂O) emissions are jeopardizing the climate goals of the Paris Agreement, according to a major new study by an international team of scientists.

The growing use of nitrogen fertilizers in the production of food worldwide is increasing atmospheric concentrations of N₂O -- a greenhouse gas 300 times more potent than carbon dioxide (CO₂) that remains in the atmosphere for more than 100 years.

Published today in the journal *Nature*, the study was led Auburn University, in the US, and involved scientists from 48 research institutions in 14 countries -- including the University of East Anglia (UEA) in the UK -- under the umbrella of the Global Carbon Project and the International Nitrogen Initiative.

The aim was to produce the most comprehensive assessment to date of all global sources and sinks of N₂O. Their findings show N₂O emissions are increasing faster than any emission scenario developed by the Intergovernmental Panel on Climate Change (IPCC), consistent with greenhouse gas scenarios that lead to global mean temperature increases well above 3°C from pre-industrial levels. The Paris Agreement aims to limit warming to less than 2°C but ideally no more than 1.5°C.

The study points to an alarming trend affecting climate change: N₂O has risen 20 per cent from pre-industrial levels -- from 270 parts per billion (ppb) in 1750 to 331ppb in 2018 -- with the fastest growth observed in the last 50 years due to emissions from human activities.

Prof Hanqin Tian, director of the International Center for Climate and Global Change Research at Auburn University's School of Forestry and Wildlife Sciences, co-led the study.

"The dominant driver of the increase in atmospheric nitrous oxide comes from agriculture, and the growing demand for food and feed for animals will further increase global nitrous oxide emissions," said Prof Tian. "There is a conflict between the way we are feeding people and stabilizing the climate."

Like CO₂, N₂O is a long-lived greenhouse gas and is also currently the most significant human-induced agent depleting the stratospheric ozone layer, which protects Earth from most of the Sun's harmful ultraviolet radiation

Lead UK author Dr Parvatha Suntharalingam, of UEA's School of Environmental Sciences, said: "This study presents the most comprehensive and detailed picture to date, of N₂O emissions and their impact on climate.

"This new analysis identifies the factors driving the steadily increasing atmospheric levels of N₂O, and highlights the urgent need to develop effective mitigation strategies if we are to limit global warming and meet climate goals."

The study presents a comprehensive global N₂O inventory that incorporates both natural and human-related sources, and accounts for the interaction between nitrogen additions to the earth system and the biochemical processes that control N₂O emissions. It covers 21 natural and human-related sectors between 1980 and 2016.

Human-induced emissions, which are dominated by nitrogen additions to croplands, increased by 30 per cent over the past four decades to 7.3 teragrams of nitrogen per year.

The analysis also reveals an emerging N₂O-climate 'feedback' resulting from interactions between nitrogen additions to crops for food production and global warming, further enhancing emissions derived from agriculture.

The study found that the largest contributors to global N₂O emissions come from East Asia, South Asia, Africa and South America. Emissions from synthetic fertilizers dominate releases in China, India and the US, while emissions from the application of livestock manure as fertilizer dominates releases in Africa and South America. The highest growth rates in emissions are in emerging economies, particularly Brazil, China and India, where crop production and livestock numbers have increased.

However, N₂O emissions in Europe decreased in agriculture and the chemical industry. This was due to a combination of factors, including voluntary measures to remove N₂O from flue gases in the Nylon industry and the introduction of an emissions trading scheme, as well as agriculture in many Western European countries moving to more efficient use of fertilizer to reduce environmental impacts such as

pollution of groundwater and surface water. Policies on nitrogen fertilizer usage were also introduced.

Study co-leader Dr Josep 'Pep' Canadell, of the Commonwealth Scientific and Industrial Research Organisation (CSIRO) in Australia, is executive director of the Global Carbon Project. He said: "This new analysis calls for a full-scale rethink in the ways we use and abuse nitrogen fertilizers globally and urges us to adopt more sustainable practices in the way we produce food, including the reduction of food waste.

"These findings underscore the urgency and opportunities to mitigate nitrous oxide emissions worldwide to avoid the worst of climate impacts."

Story Source:

Materials provided by [University of East Anglia](#). Note: Content may be edited for style and length.

Journal Reference:

1. Hanqin Tian, Rongting Xu, Josep G. Canadell, Rona L. Thompson, Wilfried Winiwarter, Parvatha Suntharalingam, Eric A. Davidson, Philippe Ciais, Robert B. Jackson, Greet Janssens-Maenhout, Michael J. Prather, Pierre Regnier, Naiqing Pan, Shufen Pan, Glen P. Peters, Hao Shi, Francesco N. Tubiello, Sönke Zaehle, Feng Zhou, Almut Arneth, Gianna Battaglia, Sarah Berthet, Laurent Bopp, Alexander F. Bouwman, Erik T. Buitenhuis, Jinfeng Chang, Martyn P. Chipperfield, Shree R. S. Dangal, Edward Dlugokencky, James W. Elkins, Bradley D. Eyre, Bojie Fu, Bradley Hall, Akihiko Ito, Fortunat Joos, Paul B. Krummel, Angela Landolfi, Goulven G. Laruelle, Ronny Lauerwald, Wei Li, Sebastian Lienert, Taylor Maavara, Michael MacLeod, Dylan B. Millet, Stefan Olin, Prabir K. Patra, Ronald G. Prinn, Peter A. Raymond, Daniel J. Ruiz, Guido R. van der Werf, Nicolas Vuichard, Junjie Wang, Ray F. Weiss, Kelley C. Wells, Chris Wilson, Jia Yang, Yuanzhi Yao. **A comprehensive quantification of global nitrous oxide sources and sinks.** *Nature*, 2020; 586 (7828): 248 DOI: [10.1038/s41586-020-2780-0](https://doi.org/10.1038/s41586-020-2780-0)

Cite This Page:

- [MLA](#)
- [APA](#)
- [Chicago](#)

University of East Anglia. "Nitrous oxide emissions pose an increasing climate threat, study finds." ScienceDaily. ScienceDaily, 7 October 2020. <www.sciencedaily.com/releases/2020/10/201007123131.htm>.